

M.H

PCT/EP 99/05744

09/622199




Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

EP 99/5744



4

EPO - DG 1

21. 10. 1999

Bescheinigung

Certificate

Attestation

(41)

Die angehefteten Unterla-  
gen stimmen mit der  
ursprünglich eingereichten  
Fassung der auf dem näch-  
sten Blatt bezeichneten  
europäischen Patentanmel-  
dung überein.

The attached documents  
are exact copies of the  
European patent application  
described on the following  
page, as originally filed.

Les documents fixés à  
cette attestation sont  
conformes à la version  
initialement déposée de  
la demande de brevet  
européen spécifiée à la  
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

98401944.8

PRIORITY DOCUMENT

Der Präsident des Europäischen Patentamts:  
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets  
p.o.

I.L.C. HATTEN-HECKMAN

DEN HAAG, DEN  
THE HAGUE,  
LA HAYE, LE

06/10/99

**THIS PAGE BLANK (USPTO)**



Europäisches  
Patentamt

European  
Patent Office

Office européen  
des brevets

**Blatt 2 der Bescheinigung**  
**Sheet 2 of the certificate**  
**Page 2 de l'attestation**

Anmeldung Nr.:  
Application no.: 98401944.8  
Demande n°:

Anmeldetag:  
Date of filing: 29/07/98  
Date de dépôt:

Anmelder:  
Applicant(s):  
Demandeur(s):  
SOCIETE CIVILE BIOPROJET  
F-75003 Paris  
FRANCE

Bezeichnung der Erfindung:  
Title of the invention:  
Titre de l'invention:

Non-imidazole aryloxy (or arylthio) alkylamines as histamine H3-receptor antagonists and their  
therapeutic applications

In Anspruch genommene Priorität(en) / Priority(ies) claimed / Priorité(s) revendiquée(s)

Staat:  
State:  
Pays:

Tag:  
Date:  
Date:

Aktenzeichen:  
File no.  
Numéro de dépôt:

Internationale Patentklassifikation:  
International Patent classification:  
Classification internationale des brevets:

C07D295/088, C07C211/08, C07D211/04, C07D295/185, C07D211/62, C07D211/70, C07D207/20, A61K31/13, A61  
K31/40, A61K31/445, A61K31/495

Am Anmeldetag benannte Vertragsstaaten:  
Contracting states designated at date of filing: AT/BE/CH/CY/DE/DK/ES/FI/FR/GB/GR/IE/IT/LI/LU/MC/NL/PT/SE  
Etats contractants désignés lors du dépôt:

Bemerkungen:  
Remarks:  
Remarques:

**THIS PAGE BLANK (USPTO)**

## NON-IMIDAZOLE ARYLOXY (OR ARYLTHIO) ALKYLAMINES AS HISTAMINE H<sub>3</sub>-RECEPTOR ANTAGONISTS AND THEIR THERAPEUTIC APPLICATIONS.

5           The present invention relates to novel aryloxy (or arylthio) alkylamines, to their preparation and to their therapeutic applications.

          Antagonists of histamine H<sub>3</sub> receptor are known especially to increase synthesis and release of cerebral histamine. Through this mechanism, they induce an extended wakefulness, an improvement in cognitive processes,  
10   a reduction in food intake and a normalization of vestibular reflexes (Schwartz et al., *Physiol. Rev.*, 1991, 71: 1-51).

          Whence these agents are potentially useful in several central nervous system disorders such as Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and  
15   motion sickness.

          All the H<sub>3</sub> receptor antagonist compounds known so far resemble histamine in possessing an imidazole ring (Ganellin et al., *Ars Pharmaceutica*, 1995, 36:3, 455-468; Stark et al., *Drug of the Future*, 1996, 21(5), 507-520).


          Nevertheless, such imidazole derivatives may show drawbacks  
20   such as poor blood-brain barrier penetration and/or some hepatic and ocular toxicities. These drawbacks, which can prevent their therapeutic development, appear to be linked to the presence of an imidazole ring substituted by a hydrophobic chain.

          Attempts to develop H<sub>3</sub> receptor antagonists without an imidazole  
25   ring have up to now been unsuccessful, the compounds being of low potency.

          In this respect, non-imidazole compounds such as betahistine (J-M. Arrang et al., *Eur. J. Pharmacol.* 1985, 111: 72-84), phencyclidine (J-M. Arrang et al., *Eur. J. Pharmacol.* 1988, 157: 31-35), dimaprit (J-C Schwartz et al., *Agents Actions* 1990, 30: 13-23), clozapine (M. Kathmann et al.,  
30   *Psychopharmacology* 1994, 116: 464-468), and sesquiterpenes (M. Takigawa et al., JP 06 345 642 (20 Dec 1994)) were suggested to display H<sub>3</sub>-receptor antagonism but all these compounds have only very low potency.

The present invention provides new compounds, the structure of which do not contain an imidazole moiety, which are useful as histamine H<sub>3</sub>-receptor antagonists while avoiding the above-mentioned drawbacks of the known H<sub>3</sub>-antagonists.

The compounds of the invention have the following general formula (I):




in which:

- C<sub>n</sub>H<sub>2n</sub> is a linear or branched hydrocarbon chain with n ranging from 2 to 8;
- X is an oxygen or sulfur atom;
- R<sup>1</sup> and R<sup>2</sup> may be identical or different and represent each independently
  - a lower alkyl or cycloalkyl,


or taken together with the nitrogen atom to which they are attached,

- a saturated nitrogen-containing ring



with m ranging from 4 to 7, or


- an unsaturated nitrogen-containing ring



with p, q and r being 1 to 3 independently,

such nitrogen-containing ring i) or ii) being unsubstituted or substituted by one or more lower alkyl or cycloalkyl, or carboalkoxy groups, or

- a morpholino group, or
- a N-substituted piperazino group:



with R being a lower alkyl, an alkanoyl or an optionally substituted phenyl group;

- $n_3$  is an integer from 0 to 5;
- $R^3$  represents each independently
  - a halogen atom,
  - a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkyloximino, aryloximino,  $\alpha$ -hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group,
  - or taken together with the carbon atoms of the phenyl ring to which it is fused, a 5- or 6-membered saturated or unsaturated ring or a benzene ring.

The invention also relates to the addition salts which the compounds form with pharmaceutically acceptable acids. The pharmaceutically acceptable salts comprise the nontoxic salt of inorganic or organic acids. Examples of these salts include the hydrochloride, the hydrobromide or the hydrogen maleate or hydrogen oxalate.


The present invention also encompasses the hydrates of the compounds, the hydrated salts of these compounds and the polymorphic crystalline structures. When the compounds can exist in one or a number of isomeric forms according to the number of asymmetric centres in the molecule, the invention relates both to all the optical isomers and to their racemic modifications and the corresponding diastereoisomers. The separation of the diastereoisomers and/or of the optical isomers can be carried out according to methods known per se.

According to the invention, lower alkyl or cycloalkyl is intended to mean a linear or branched alkyl group containing from 1 to 6 carbon atoms, or a saturated carbocycle containing 3 to 6 carbon atoms.

Typically examples of lower alkyl are methyl, ethyl, propyl, isopropyl and butyl groups.


A preferred group of compounds according to the invention comprises those with  $R^1$  and  $R^2$  representing independently a lower alkyl group, especially an ethyl group.

Preferred compounds are also those of formula (I) in which  $R^1$  and  $R^2$  taken together with the nitrogen atom to which they are attached, form a saturated nitrogen-containing ring:



especially with  $m$  being 4, 5 or 6, optionally substituted with an alkyl group, preferably a methyl group.

10                      Another preferred group of compounds comprises compounds (I) in which  $R^1$  and  $R^2$  taken together with the nitrogen atom to which they are attached, form an unsaturated nitrogen-containing ring:



especially with  $p$ ,  $q$ , and  $r$  being 1 or 2. In this group, more preferred compounds are those with  $p$  being 2 and  $q$  and  $r$  each being 1.

20                      Typical example of  $-\text{NR}^1\text{R}^2$  representing a N-substituted piperazino group is N-acetylpiperazino.

A preferred group of compounds according to the invention is the group composed of compounds of formula (I) in which  $X$  is an oxygen atom.

25                      Another preferred group of compounds comprises compounds (I) in which  $-\text{C}_n\text{H}_{2n}-$  is a linear chain  $-(\text{CH}_2)_n-$  with  $n$  being as previously defined.

Preferred compounds are also those with  $n$  varying from 3 to 5, and with  $n$  being more preferably 3.

30                      A sub-class of compounds according to the invention comprises the compounds of formula (I) with  $n_3$  being zero that is those having an unsubstituted phenyl moiety.

Another group of compounds according to the invention is composed of compounds containing one or more substituents  $R^3$  which may be identical or different. In this group, the compounds having a mono- or di-



substituted ( $n_3 = 1$  or  $2$ ) phenyl moiety are preferred and those mono-substituted with one group  $R^3$  as defined above in para-position are particularly preferred.

Among these compounds, ( $n_3$  being  $1$ )  $R^3$  is preferably a halogen atom or a cyano, nitro, alkanoyl, alkyloximino or  $\alpha$ -hydroxyalkyl group.

Still more preferred compounds are those with  $R^3$  being CN, NO<sub>2</sub>, COCH<sub>3</sub>, COC<sub>2</sub>H<sub>5</sub>, H<sub>3</sub>C-C=N-OH, H<sub>3</sub>C-CH-OH.

$R^3$  being a halogen atom may be advantageously selected from fluorine, chlorine and bromine.

$R^3$  being an aryl group, may be especially a phenyl group.

In the other substituents  $R^3$ , the aryl moiety is advantageously a phenyl moiety.

$R^3$  being an aryloxy group may be especially a phenoxy group.

According to the invention, alkanoyl is intended to mean a group containing an alkyl moiety as defined above.

Typical examples of  $R^3$  being an alkanoyl, aroyl or arylalkanoyl group are acetyl, butyryl and propionyl groups, benzoyl group or phenylacetyl group.

Typical examples of  $R^3$  forming together with the carbon atoms of the phenyl ring to which it is fused, a saturated ring leads to 5,6,7,8-tetrahydronaphthyl or forming a benzene ring leads to a naphthyl moiety.

According to the invention, alkenyl or alkynyl group may contain advantageously from 1 to 8 carbon atoms, in particular from 1 to 6 carbon atoms and preferably 1 to 4 carbon atoms.

In carboalkoxy, carboxyamido or carboxamide groups, the hydrocarbon chain is saturated, linear or branched and contains an alkyl moiety as defined above.

In alkoxy, alkyloximino, arylalkyl or  $\alpha$ -hydroxyalkyl group, the alkyl moiety is as previously defined also.

Particularly preferred compounds are:

1-(5-phenoxypropyl)-piperidine

1-(5-phenoxypropyl)-pyrrolidine

N-methyl-N-(5-phenoxypropyl)-ethylamine

- 1-(5-phenoxypropyl)-morpholine  
N-(5-phenoxypropyl)-hexamethyleneimine  
N-ethyl-N-(5-phenoxypropyl)-propylamine  
1-(5-phenoxypropyl)-2-methyl-piperidine  
5 1-(5-phenoxypropyl)-4-propyl-piperidine  
1-(5-phenoxypropyl)-4-methyl-piperidine  
1-(5-phenoxypropyl)-3-methyl-piperidine  
1-acetyl-4-(5-phenoxypropyl)-piperazine  
1-(5-phenoxypropyl)-3,5-trans-dimethyl-piperidine  
10 1-(5-phenoxypropyl)-3,5-cis-dimethyl-piperidine  
1-(5-phenoxypropyl)-2,6-cis-dimethyl-piperidine  
4-carboethoxy-1-(5-phenoxypropyl)-piperidine  
3-carboethoxy-1-(5-phenoxypropyl)-piperidine  
1-(5-phenoxypropyl)-1,2,3,6-tetrahydropyridine  
15 1-[5-(4-nitrophenoxy)-propyl]-pyrrolidine  
1-[5-(4-chlorophenoxy)-propyl]-pyrrolidine  
1-[5-(4-methoxyphenoxy)-propyl]-pyrrolidine  
1-[5-(4-methylphenoxy)-propyl]-pyrrolidine  
1-[5-(4-cyanophenoxy)-propyl]-pyrrolidine  
20 1-[5-(2-naphthoxy)-propyl]-pyrrolidine  
1-[5-(1-naphthoxy)-propyl]-pyrrolidine  
1-[5-(3-chlorophenoxy)-propyl]-pyrrolidine  
1-[5-(4-phenylphenoxy)-propyl]-pyrrolidine  
1-[5-{2-(5,6,7,8-tetrahydronaphthyl)-oxy}-propyl]-pyrrolidine  
25 1-[5-(3-phenylphenoxy)-propyl]-pyrrolidine  
1-(5-phenoxypropyl)-2,5-dihydropyrrole  
1-[5-{1-(5,6,7,8-tetrahydronaphthyl)-oxy}-propyl]-pyrrolidine  
1-(4-phenoxybutyl)-pyrrolidine  
1-(6-phenoxyhexyl)-pyrrolidine  
30 1-(5-phenylthiopropyl)-pyrrolidine  
1-(4-phenylthiobutyl)-pyrrolidine  
1-(3-phenoxypropyl)-pyrrolidine  
1-[5-(3-nitrophenoxy)-propyl]-pyrrolidine

- 1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine  
1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine  
1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine  
1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine  
5 1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine  
N-[3-(4-nitrophenoxy)-propyl]-diethylamine  
N-[3-(4-cyanophenoxy)-propyl]-diethylamine  
1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine  
1-[5-[4-(phenylacetyl)-phenoxy]-pentyl]-pyrrolidine  
10 N-[3-(4-acetylphenoxy)-propyl]-diethylamine  
1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine  
1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine  
1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine  
1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]-pyrrolidine  
15 1-[5-(4-cyanophenoxy)-pentyl]-diethylamine  
1-[5-(4-cyanophenoxy)-pentyl]-piperidine  
N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine  
N-[2-(4-cyanophenoxy)-ethyl]-diethylamine  
N-[3-(4-cyanophenoxy)-propyl]-dimethylamine  
20 N-[4-(4-cyanophenoxy)-butyl]-diethylamine  
N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine  
1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine  
1-[3-(4-cyanophenoxy)-propyl]-piperidine  
N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine  
25 N-[6-(4-cyanophenoxy)-hexyl]-diethylamine  
N-[3-(4-cyanophenoxy)-propyl]-dipropylamine  
N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine  
4-(3-diethylaminopropoxy)-acetophenone-oxime  
1-[3-(4-acetylphenoxy)-propyl]-piperidine  
30 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine  
1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-piperidine  
1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine  
1-[3-(4-propionylphenoxy)-propyl]-piperidine

1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine

1-[3-(4-formylphenoxy)-propyl]-piperidine

1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine

N-[3-(4-propionylphenoxy)-propyl]-diethylamine

5 1-[3-(4-butyrylphenoxy)-propyl]-piperidine

1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine

More preferred compounds are:

1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine

N-[3-(4-cyanophenoxy)-propyl]-diethylamine

10 N-[3-(4-acetylphenoxy)-propyl]-diethylamine

1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]-pyrrolidine

N-[4-(4-cyanophenoxy)-butyl]-diethylamine

1-[3-(4-cyanophenoxy)-propyl]-piperidine

N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine

15 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine

4-(3-diethylaminopropoxy)-acetophenone-oxime

1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine

1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine

1-[3-(4-propionylphenoxy)-propyl]-piperidine

20 Compounds of formula (I) in which:

- $-NR^1R^2$  is a pyrrolidinyl group,  $C_nH_{2n}$  is a linear chain  $-(CH_2)_n-$  and  $n_3$  is zero, X being an oxygen atom with n ranging from 3 to 5, or X being a sulfur atom with n being 4 or 5;
- $-NR^1R^2$  is a piperidinyl group,  $C_nH_{2n}$  is a linear chain  $-(CH_2)_n-$  and X is an oxygen atom,  $n_3$  being zero with n being 2, 5 or 8 or  $n_3$  being 1 with  $R^3$  being 4-CN and n being 5;
- $-NR^1R^2$  is a diethylamine group, X is an oxygen atom,  $C_nH_{2n}$  is a linear chain  $-(CH_2)_n-$  and  $n_3$  is 1,  $R^3$  being 4- $NO_2$  or 4- $COCH_3$  with n being 3 or  $R^3$  being 4-CN with n being 2 to 4;
- $-NR^1R^2$  is a dimethylamine group, X is an oxygen atom,  $C_nH_{2n}$  is a linear chain  $-(CH_2)_n-$  and  $n_3$  is 1,  $R^3$  being 4-CN with n being 3,


30

are known in the art.


A subject of the invention is thus the use of these compounds as antagonists at the histamine H<sub>3</sub>-receptors, in particular to prepare medicaments acting as H<sub>3</sub>-antagonists intended for the treatments detailed below.

The compounds according to the invention may be prepared  
5 according to one of the following schemes 1-5:


SCHEME 1 (methods A, B, C, D, H and K):




SCHEME 2 (methods F and L):




SCHEME 3 (method E):



SCHEME 4 (method G):



SCHEME 5 (method J):



In these schemes,  $R^1$ ,  $R^2$ ,  $R^3$ , X and n are as defined in general formula (I).

Me and Et are intended to mean methyl and ethyl.

Detailed synthesis procedures are given in the examples.

5        The compounds of formula (I) according to the invention have antagonistic properties at the histamine  $H_3$ -receptors. They cause an increase in synthesis and release of cerebral histamine.

This property makes the compounds of the invention useful derivatives in human or veterinary medicine.

10        Their therapeutical applications are those known for  $H_3$ -antagonist compounds and especially relate to the central nervous system disorders such as Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and motion sickness.

15        Therefore, the compounds of formula (I) according to the invention are advantageously used as active ingredient of medicaments which act as an antagonist of  $H_3$ -receptors of histamine, in particular of medicaments having psychotropic effects, promoting wakefulness, attention, memory and improving mood, in treatment of pathologies such as Alzheimer disease and other cognitive disorders in aged persons, depressive or simply asthenic states.

20        Their nootropic effects can be useful to stimulate attention and memorization capacity in healthy humans.

In addition, these agents can be useful in treatment of obesity, vertigo and motion sickness.

25        It can also be useful to associate the compounds of the invention with other psychiatric agents such as neuroleptics to increase their efficiency and reduce their side effects.

Application in certain form of epilepsy is also foreseen.

Their therapeutic applications involve also peripheral organs mainly a stimulant of secretions or gastro-intestinal motricity.

30        The compounds of the invention are particularly useful for the treatment of CNS disorders of aged persons.

The present invention also relates to medicaments having the above-mentioned effects comprising as active ingredient, a therapeutically effective amount of a compound of formula (I).

5 The present invention also relates to pharmaceutical compositions containing as active ingredient, a therapeutically effective amount of a compound (I) together with a pharmaceutically acceptable vehicle or excipient.

The medicaments or pharmaceutical compositions according to the invention can be administered via oral, parenteral or topical routes, the active ingredient being combined with a therapeutically suitable excipient or  
10 vehicle.

According to the invention, oral administration is advantageously used.

Another subject of the present invention is the use of the compounds of formula (I) for the preparation of H<sub>3</sub>-antagonist medicaments  
15 according to the above-mentioned forms.

The invention further relates to the use of the compounds of formula (I) for preparing medicaments having the pre-cited effects.

Still another subject of the invention is a method for the treatment of precited ailments comprising administering a therapeutically effective dose of  
20 a compound (I), optionally in combination with a therapeutically acceptable vehicle or excipient.

For each of the above-indications, the amount of the active ingredient will depend upon the condition of the patient.

However, a suitable effective dose will be in general in the range  
25 of from 10 to 500 mg per day and of from 1 to 10 mg/day for particularly active compounds.

These doses are given on the basis of the compound and should be adapted for the salts, hydrates or hydrated salts thereof.

The invention is now illustrated by the following examples.






30






### EXAMPLES

The structure of the synthesized compounds and their method of preparation as well as their melting point, recrystallisation solvent and elemental analysis are summarized in the following Table I:














TABLE 1:






Nº	FORMULA STRUCTURE NAME	mp (recryst. solv)	analysis (calc.)	method
1	$C_{16}H_{25}NO; C_2H_2O_4$  1-(5-phenoxypropyl)-piperidine hydrogen oxalate	143-145°C (absolute ethanol)	C: 64.06 (64.07) H: 8.09 (8.16) N: 4.14 (4.15)	A
2	$C_{15}H_{23}NO; C_2H_2O_4$  1-(5-phenoxypropyl)-pyrrolidine hydrogen oxalate	153-155°C (absolute ethanol)	C: 63.06 (63.14) H: 7.78 (7.79) N: 4.42 (4.33)	A
3	$C_{14}H_{23}NO; C_2H_2O_4$  N-methyl-N-(5-phenoxypropyl)-ethylamine hydrogen oxalate	122-124°C (absolute ethanol)	C: 61.74 (61.72) H: 8.24 (8.09) N: 4.52 (4.50)	A
4	$C_{15}H_{23}NO_2; C_2H_2O_4$  1-(5-phenoxypropyl)-morpholine hydrogen oxalate	166-168°C (absolute ethanol)	C: 60.10 (60.16) H: 7.45 (7.31) N: 4.08 (4.13)	A
5	$C_{17}H_{27}NO; C_2H_2O_4$  N-(5-phenoxypropyl)-hexamethyleneimine hydrogen oxalate	132-134°C (absolute ethanol)	C: 64.70 (64.93) H: 8.34 (8.32) N: 3.85 (3.99)	A







6	$C_{16}H_{27}NO$ ; $C_2H_2O_4$  N-ethyl-N-(5-phenoxypropyl)-propylamine hydrogen oxalate	90-91°C (isopropyl alcohol)	C: 63.60 (63.69) H: 8.81 (8.61) N: 3.97 (4.13)	B
7	$C_{17}H_{27}NO$ ; 1.1 $C_2H_2O_4$  1-(5-phenoxypropyl)-2-methyl-piperidine hydrogen oxalate	80-83°C (isopropyl alcohol)	C: 64.15 (63.98) H: 8.42 (8.17) N: 3.97 (3.89)	B
8	$C_{19}H_{31}NO$ ; $C_2H_2O_4$  1-(5-phenoxypropyl)-4-propyl-piperidine hydrogen oxalate	165-166°C (absolute ethanol)	C: 66.27 (66.46) H: 8.94 (8.76) N: 3.72 (3.69)	B
9	$C_{17}H_{27}NO$ ; $C_2H_2O_4$  1-(5-phenoxypropyl)-4-methyl-piperidine hydrogen oxalate	151-152°C (absolute ethanol)	C: 64.87 (64.93) H: 8.41 (8.32) N: 4.01 (3.99)	B
10	$C_{17}H_{27}NO$ ; $C_2H_2O_4$  1-(5-phenoxypropyl)-3-methyl-piperidine hydrogen oxalate	140-141°C (isopropyl alcohol)	C: 65.35 (64.93) H: 8.49 (8.32) N: 4.00 (3.99)	B






11	$C_{17}H_{25}N_2O_2; C_2H_2O_4$  1-acetyl-4-(5-phenoxy-pentyl)-piperazine hydrogen oxalate	186-188°C (absolute ethanol)	C: 59.78 (59.99) H: 7.47 (7.42) N: 7.35 (7.36)	B
12	$C_{18}H_{29}NO; 1.05 C_2H_2O_4$  1-(5-phenoxy-pentyl)-3,5-trans-dimethyl- piperidine hydrogen oxalate	154-155°C (absolute ethanol)	C: 65.16 (65.25) H: 8.61 (8.47) N: 3.66 (3.79)	B
13	$C_{18}H_{29}NO; C_2H_2O_4$  1-(5-phenoxy-pentyl)-3,5-cis-dimethyl- piperidine hydrogen oxalate	154-155°C (isopropyl alcohol)	C: 65.62 (65.73) H: 8.64 (8.55) N: 3.63 (3.83)	B
14	$C_{18}H_{29}NO; HCl$  1-(5-phenoxy-pentyl)-2,6-cis-dimethyl- piperidine hydrochloride	135-136°C (acetone)	C: 69.18 (69.32) H: 9.79 (9.70) N: 4.28 (4.49)	B
15	$C_{19}H_{29}NO_3; C_2H_2O_4$  4-carboethoxy-1-(5-phenoxy-pentyl)- piperidine hydrogen oxalate	149-150°C (absolute ethanol)	C: 61.16 (61.60) H: 7.76 (7.63) N: 3.40 (3.42)	B

16	$C_{19}H_{29}NO_3; C_2H_2O_4$  3-carboethoxy-1-(5-phenoxy)pentyl-piperidine hydrogen oxalate	117-118°C (isopropyl alcohol)	C: 61.54 (61.60) H: 7.87 (7.63) N: 3.29 (3.42)	B
17	$C_{16}H_{23}NO; C_2H_2O_4$  1-(5-phenoxy)pentyl-1,2,3,6-tetrahydropyridine hydrogen oxalate	177-179°C (methanol)	C: 64.19 (64.46) H: 7.49 (7.51) N: 4.25 (4.18)	B
18	$C_{15}H_{22}N_2O_3; C_2H_2O_4; 0.2 H_2O$  1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	145-147°C (absolute ethanol)	C: 54.89 (54.89) H: 6.68 (6.61) N: 7.41 (7.53)	C
19	$C_{15}H_{22}ClNO; C_2H_2O_4$  1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	139-141°C (absolute ethanol)	C: 57.00 (57.06) H: 6.63 (6.76) N: 3.79 (3.91) Cl: 10.24 (9.91)	C
20	$C_{16}H_{25}NO_2; C_2H_2O_4$  1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	115-116°C (absolute ethanol)	C: 61.22 (61.17) H: 7.72 (7.70) N: 4.03 (3.96)	C
21	$C_{16}H_{25}NO; C_2H_2O_4$  1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	138-140°C (absolute ethanol)	C: 64.05 (64.07) H: 8.00 (8.07) N: 4.10 (4.15)	C






22	$C_{16}H_{22}N_2O$ ; 1.1 $C_2H_2O_4$  1.1 $(COOH)_2$ 1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	129-130°C (absolute ethanol)	C: 61.24 (61.16) H: 6.81 (6.82) N: 7.95 (7.84)	C
23	$C_{19}H_{25}NO$ ; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(2-naphthyloxy)-pentyl]-pyrrolidine hydrogen oxalate	166-167°C (methanol)	C: 67.42 (67.54) H: 7.26 (7.29) N: 3.66 (3.75)	C
24	$C_{19}H_{25}NO$ ; 1.25 $C_2H_2O_4$  1.25 $(COOH)_2$ 1-[5-(1-naphthyloxy)-pentyl]-pyrrolidine hydrogen oxalate	160-163°C (methanol)	C: 65.12 (65.22) H: 7.17 (7.00) N: 3.52 (3.54)	C
25	$C_{15}H_{22}ClNO$ ; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	131-132°C (absolute ethanol)	C: 56.94 (57.06) H: 6.67 (6.76) N: 3.74 (3.91) Cl: 9.64 (9.91)	C
26	$C_{21}H_{27}NO$ ; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	189-190°C (methanol)	C: 69.16 (69.15) H: 7.39 (7.32) N: 3.39 (3.51)	C






27	$C_{19}H_{29}NO$ ; $C_2H_2O_4$  1-[5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine hydrogen oxalate	131-132°C (absolute ethanol)	C: 66.73 (66.82) H: 8.37 (8.28) N: 3.68 (3.71)	C
28	$C_{21}H_{27}NO$ ; 1.1 $C_2H_2O_4$  1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	155-157°C (absolute ethanol)	C: 68.40 (68.22) H: 7.04 (7.21) N: 3.45 (3.43)	C
29	$C_{15}H_{21}NO$ ; $C_2H_2O_4$  1-(5-phenoxy)pentyl-2,5-dihydropyrrole hydrogen oxalate	140-141°C (absolute ethanol)	C: 63.45 (63.54) H: 7.26 (7.21) N: 4.26 (4.36)	B
30	$C_{19}H_{29}NO$ ; $C_2H_2O_4$  1-[5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine hydrogen oxalate	148-149°C (absolute ethanol)	C: 66.99 (66.82) H: 8.47 (8.28) N: 3.72 (3.71)	C
31	$C_{14}H_{21}NO$ ; $C_2H_2O_4$  1-(4-phenoxybutyl)-pyrrolidine hydrogen oxalate	143-144°C (absolute ethanol)	C: 62.25 (62.12) H: 7.46 (7.49) N: 4.49 (4.53)	C






32	$C_{16}H_{25}NO$ ; 1.1 $C_2H_2O_4$  1.1 $(COOH)_2$ 1-(6-phenoxyhexyl)-pyrrolidine hydrogen oxalate	146-147°C (absolute ethanol)	C: 63.06 (63.10) H: 8.03 (7.91) N: 4.32 (4.04)	C
33	$C_{15}H_{23}NS$ ; 1.1 $C_2H_2O_4$  1.1 $(COOH)_2$ 1-(5-phenylthiopentyl)-pyrrolidine hydrogen oxalate	150-152°C (absolute ethanol)	C: 59.52 (59.29) H: 7.44 (7.29) N: 4.06 (4.02)	C
34	$C_{14}H_{21}NS$ ; $C_2H_2O_4$  $(COOH)_2$ 1-(4-phenylthiobutyl)-pyrrolidine hydrogen oxalate	114-116°C (absolute ethanol)	C: 59.24 (59.05) H: 7.16 (7.12) N: 4.16 (4.30) S: 9.79 (9.85)	C
35	$C_{13}H_{19}NO$ ; $C_2H_2O_4$  $(COOH)_2$ 1-(3-phenoxypropyl)-pyrrolidine hydrogen oxalate	169-170°C (absolute ethanol)	C: 60.98 (61.00) H: 7.14 (7.17) N: 4.64 (4.74)	C
36	$C_{15}H_{22}N_2O_3$ ; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	130-131°C (absolute ethanol)	C: 55.30 (55.43) H: 6.55 (6.57) N: 7.49 (7.60)	C
37	$C_{15}H_{22}FNO$ ; $C_2H_2O_4$  $(COOH)_2$ 1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	149-150°C (absolute ethanol)	C: 59.52 (59.81) H: 7.12 (7.09) N: 4.05 (4.10)	C







38	$C_{17}H_{26}N_2O_3$ ; $C_2H_2O_4$  1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine hydrogen oxalate	148-149°C (absolute ethanol)	C: 57.32 (57.55) H: 7.19 (7.12) N: 6.89 (7.07)	C
39	$C_{17}H_{25}NO_2$ ; $C_2H_2O_4$  1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	130-134°C (absolute ethanol)	C: 62.43 (62.45) H: 7.41 (7.45) N: 3.75 (3.83)	D
40	$C_{15}H_{24}N_2O$ ; 2.1 $C_2H_2O_4$  1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine di-(hydrogen oxalate)	120-122°C (absolute ethanol)	C: 52.49 (52.72) H: 6.74 (6.50) N: 6.32 (6.40)	E <sub>1</sub>
41	$C_{16}H_{22}N_2O$ ; $C_2H_2O_4$  1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	119-120°C (absolute ethanol)	C: 61.95 (62.05) H: 6.88 (6.94) N: 8.00 (8.04)	C
42	$C_{13}H_{20}N_2O_3$ ; $C_2H_2O_4$  N-[3-(4-nitrophenoxy)-propyl]-diethylamine hydrogen oxalate	160-161°C (absolute ethanol/ methanol 1:1)	C: 52.46 (52.63) H: 6.49 (6.48) N: 8.10 (8.12)	F













43	$C_{14}H_{20}N_2O; C_2H_2O_4$  N-[3-(4-cyanophenoxy)-propyl]-diethylamine hydrogen oxalate	148-150°C (absolute ethanol)	C: 59.40 (59.62) H: 6.82 (6.88) N: 8.60 (8.69)	F
44	$C_{22}H_{27}NO_2; C_2H_2O_4$  1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	141-142°C (absolute ethanol)	C: 67.17 (67.43) H: 6.80 (6.84) N: 3.18 (3.28)	D
45	$C_{23}H_{29}NO_2; C_2H_2O_4$  1-[5-[4-(phenylacetyl)-phenoxy]-pentyl]-pyrrolidine hydrogen oxalate	177-178°C (absolute ethanol)	C: 67.77 (68.01) H: 7.09 (7.08) N: 3.26 (3.17)	D
46	$C_{15}H_{23}NO_2; 1.1 C_2H_2O_4$  N-[3-(4-acetylphenoxy)-propyl]-diethylamine hydrogen oxalate	108-110°C (absolute ethanol)	C: 59.30 (59.30) H: 7.47 (7.29) N: 4.18 (4.02)	F
47	$C_{17}H_{26}N_2O_2; C_2H_2O_4$  1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	142-144°C (absolute ethanol)	C: 59.67 (59.99) H: 7.55 (7.42) N: 7.25 (7.36)	C



48	$C_{21}H_{27}NO_2$ ; $C_2H_2O_4$  1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine hydrogen oxalate	135-136°C (absolute ethanol)	C: 66.49 (66.49) H: 7.05 (7.04) N: 3.24 (3.37)	D
49	$C_{22}H_{28}N_2O_2$ ; 1.1 $C_2H_2O_4$  1.1 (COOH) <sub>2</sub> 1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine hydrogen oxalate	176-178°C (absolute ethanol)	C: 64.56 (64.38) H: 6.89 (6.74) N: 6.26 (6.20)	E <sub>2</sub>
50	$C_{17}H_{27}NO_2$ ; $C_2H_2O_4$  1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]- pyrrolidine hydrogen oxalate	102-104°C (absolute ethanol)	C: 61.89 (62.11) H: 7.94 (7.96) N: 3.77 (3.81)	G
51	$C_{16}H_{24}N_2O$ ; $C_2H_2O_4$  N-[5-(4-cyanophenoxy)-pentyl]-diethylamine hydrogen oxalate	120-122°C (absolute ethanol)	C: 61.56 (61.70) H: 7.54 (7.48) N: 7.87 (7.99)	H
52	$C_{17}H_{24}N_2O$ ; $C_2H_2O_4$  1-[5-(4-cyanophenoxy)-pentyl]-piperidine hydrogen oxalate	115-116°C (absolute ethanol)	C: 62.62 (62.97) H: 7.20 (7.23) N: 7.76 (7.73)	H

53	$C_{14}H_{20}N_2O; C_2H_2O_4$  N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine hydrogen oxalate	148-149°C (absolute ethanol)	C: 59.68 (59.62) H: 6.76 (6.88) N: 8.57 (8.69)	H
54	$C_{13}H_{18}N_2O; C_2H_2O_4$  N-[2-(4-cyanophenoxy)-ethyl]-diethylamine hydrogen oxalate	124-125°C (absolute ethanol)	C: 58.15 (58.43) H: 6.30 (6.54) N: 8.95 (9.09)	H
55	$C_{12}H_{16}N_2O; C_2H_2O_4$  N-[3-(4-cyanophenoxy)-propyl]-dimethylamine hydrogen oxalate	166-167°C (absolute ethanol/ methanol 1:1)	C: 57.01 (57.14) H: 6.02 (6.16) N: 9.46 (9.52)	H
56	$C_{15}H_{22}N_2O; C_2H_2O_4$  N-[4-(4-cyanophenoxy)-butyl]-diethylamine hydrogen oxalate	143-145°C (absolute ethanol)	C: 60.80 (60.70) H: 7.11 (7.19) N: 8.22 (8.33)	H
57	$C_{18}H_{28}N_2O; C_2H_2O_4$  N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine hydrogen oxalate	134-136°C (absolute ethanol)	C: 63.38 (63.47) H: 8.11 (7.99) N: 7.29 (7.40)	H

58	$C_{14}H_{18}N_2O$ ; 1.1 $C_2H_2O_4$  1.1 (COOH) <sub>2</sub> 1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine hydrogen oxalate	163-165°C (absolute ethanol)	C: 58.95 (59.08) H: 6.23 (6.18) N: 8.43 (8.51)	H
59	$C_{15}H_{20}N_2O$ ; 1.05 $C_2H_2O_4$  1.05 (COOH) <sub>2</sub> 1-[3-(4-cyanophenoxy)-propyl]-piperidine hydrogen oxalate	151-153°C (absolute ethanol)	C: 60.62 (60.61) H: 6.66 (6.57) N: 8.25 (8.27)	H
60	$C_{16}H_{22}N_2O$ ; 1.05 $C_2H_2O_4$  1.05 (COOH) <sub>2</sub> N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine hydrogen oxalate	124-125°C (absolute ethanol)	C: 61.62 (61.60) H: 6.94 (6.88) N: 7.87 (7.94)	H
61	$C_{17}H_{26}N_2O$ ; $C_2H_2O_4$  (COOH) <sub>2</sub> N-[6-(4-cyanophenoxy)-hexyl]-diethylamine hydrogen oxalate	110-112°C (absolute ethanol)	C: 62.90 (62.62) H: 7.76 (7.74) N: 7.61 (7.69)	H
62	$C_{16}H_{24}N_2O$ ; $C_2H_2O_4$  (COOH) <sub>2</sub> N-[3-(4-cyanophenoxy)-propyl]-dipropylamine hydrogen oxalate	127-128°C (absolute ethanol)	C: 61.57 (61.70) H: 7.57 (7.48) N: 7.91 (7.99)	H
63	$C_{15}H_{25}NO_2$ ; $C_2H_2O_4$ ; 0.5 $H_2O$  (COOH) <sub>2</sub> 0.5 $H_2O$ N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine hydrogen oxalate hemihydrate	33-36°C (isopropyl alcohol)	C: 58.15 (58.27) H: 8.15 (8.05) N: 4.21 (4.00)	G

64	$C_{15}H_{24}N_2O_2$ ; $C_2H_2O_4$  4'-(3-diethylaminopropoxy)-acetophenone-oxime hydrogen oxalate	99-100°C (absolute ethanol)	C: 57.26 (57.61) H: 7.47 (7.39) N: 7.72 (7.90)	J
65	$C_{16}H_{23}NO_2$ ; $C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-piperidine hydrogen oxalate	159-160°C (absolute ethanol)	C: 61.18 (61.52) H: 7.11 (7.17) N: 3.96 (3.99)	K
66	$C_{17}H_{25}NO_2$ ; $C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine hydrogen oxalate	143-144°C (absolute ethanol)	C: 62.11 (62.45) H: 7.41 (7.45) N: 3.79 (3.83)	K
67	$C_{18}H_{27}NO_2$ ; $C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl- piperidine hydrogen oxalate	171-172°C (absolute ethanol)	C: 63.06 (63.31) H: 7.44 (7.70) N: 3.64 (3.69)	K
68	$C_{17}H_{25}NO_2$ ; $C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine hydrogen oxalate	160-161°C (absolute ethanol)	C: 62.47 (62.45) H: 7.46 (7.45) N: 3.77 (3.83)	K

69	$C_{17}H_{25}NO_2; C_2H_2O_4$  1-[3-(4-propionylphenoxy)-propyl]-piperidine hydrogen oxalate	148-149°C (absolute ethanol)	C: 62.54 (62.45) H: 7.51 (7.45) N: 3.79 (3.83)	L
70	$C_{18}H_{27}NO_2; C_2H_2O_4$  1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl- piperidine hydrogen oxalate	174-175°C (absolute ethanol)	C: 63.22 (63.31) H: 7.60 (7.70) N: 3.64 (3.69)	K
71	$C_{15}H_{21}NO_2; C_2H_2O_4$  1-[3-(4-formylphenoxy)-propyl]-piperidine hydrogen oxalate	152-153°C (absolute ethanol)	C: 60.23 (60.52) H: 6.81 (6.87) N: 4.15 (4.15)	L
72	$C_{18}H_{27}NO_2; C_2H_2O_4$  1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine hydrogen oxalate	121-122°C (absolute ethanol)	C: 63.02 (63.31) H: 7.73 (7.70) N: 3.66 (3.69)	L
73	$C_{16}H_{25}NO_2; 1.5 C_2H_2O_4$  N-[3-(4-propionylphenoxy)-propyl]-diethylamine hydrogen oxalate	118-120°C (absolute ethanol)	C: 57.27 (57.28) H: 7.00 (7.08) N: 3.47 (3.52)	L

74	$C_{18}H_{27}NO_2; C_2H_2O_4$  1-[3-(4-butyrylphenoxy)-propyl]-piperidine hydrogen oxalate	138-139°C (absolute ethanol)	C: 63.09 (63.31) H: 7.78 (7.70) N: 3.75 (3.69)	L
75	$C_{16}H_{21}NO_2; 1.1 C_2H_2O_4$  1.1 (COOH) <sub>2</sub> 1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6- tetrahydropyridine hydrogen oxalate	143-144°C (absolute ethanol)	C: 61.21 (61.00) H: 6.25 (6.52) N: 4.00 (3.91)	K

Compounds 1 to 75 are prepared according to the following procedures:

**METHOD A:**

A solution of 1-bromo-5-phenoxy pentane (1.4 to 3.5 mmol) in ten equivalents of the suitable secondary amine was heated to reflux temperature with stirring for 48 hours (compds. 1, 3 and 4), 24 hours (compd. 2) or 4 hours (compd. 5). After cooling, the excess base was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate formed was washed with diethyl ether and recrystallised from absolute ethanol.

**METHOD B:**

A solution of 1-bromo-5-phenoxy pentane (0.9 to 1.7 mmol) and an excess of the suitable secondary amine (2.3 to 10 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 48 hours (compd. 6) or 24 hours (compds. 7, 8, 9, 10, 11, 12&13, 14, 15, 16, 17 and 29). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The cis and trans isomers 12 and 13 were separated by column chromatography on silica gel eluting with a solvent mixture of petroleum spirit (bp 60-80°C), diethyl ether and triethylamine in the ratio 66:33:1, and the eluent was removed under reduced pressure to leave an oil. Compounds 14 and 16 were purified by column chromatography on silica gel eluting with diethyl ether and triethylamine in the ratio 99:1, and the eluent was removed under reduced pressure to leave an oil. The oil was converted to oxalate salt (compds. 6, 7, 8, 9, 11, 12, 13, 15, 16, 17 and 29) by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents of oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was



added to form a precipitate. The solid was washed with diethyl ether and recrystallised from isopropyl alcohol (compds. 6, 7, 10, 13 and 16), absolute ethanol (compds. 8, 9, 11, 12, 15 and 29) or methanol (compd. 17). The oil was converted to hydrochloride salt (compd. 14) by adding 2N HCl. The precipitate  
5 was formed in a mixture of chloroform and diethyl ether (1:1) and recrystallised from acetone.

#### METHOD C:

A solution of the suitable  $\alpha$ -bromo- $\omega$ -aryloxy alkane (0.4 to 1.4 mmol) or  $\omega$ -bromoalkyl phenyl sulphide (1 mmol, compds. 33 and 34) and an excess of  
10 pyrrolidine (10 to 15 equivalents) or 3-methylpiperidine (10 equivalents, compd. 38) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 24 hours or 16 hours (compd. 47). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The  
15 product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was  
20 added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

#### METHOD D:

A solution of the suitable 4'-(5-bromopentoxy)phenyl ketone (0.7 to 1 mmol, compds. 39, 44 and 45) or 1-bromo, 5-(4-phenoxyphenoxy)pentane (0.6 mmol, compd. 48) and an excess of pyrrolidine (10 to 15 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 16 hours (compds. 39, 44 and 48) or 24 hours (compd. 45). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium  
30 hydroxide. The product was extracted with chloroform (compds. 39, 45 and 48) or dichloromethane (compd. 44), the organic extracts dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute

ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate was washed with diethyl ether and recrystallised from absolute ethanol (recrystallised twice from absolute ethanol in the case of compd. 39).

5 **METHOD E:**

1. The oxalate 18 was prepared according to method C. A solution of compound 18 (0.57 mmol) in 10 ml methanol and 10 ml absolute ethanol was placed with 100 mg of palladium (5%) on carbon catalyst in a two-neck round-bottom flask fitted with a balloon filled with hydrogen. The mixture was stirred  
10 vigorously at room temperature and the flask was purged of air and filled with hydrogen. After 3 hours, the catalyst was filtered off on celite and the solvent removed under reduced pressure. The residual solid was converted to oxalate salt by dissolving in methanol and adding a solution of oxalic acid (2 equivalents) in absolute ethanol. Diethyl ether was added to form a precipitate.  
15 The product was recrystallised from absolute ethanol.

2. To a solution of compound 40 (0.35 mmol) in pyridine vigorously stirred at 0°C was added dropwise a slight excess of benzoyl chloride (0.4 mmol). The stirring was allowed to continue 20 minutes after the end of the addition after which the mixture was placed in the refrigerator overnight (16  
20 hours). The solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with chloroform, the organic extracts dried over magnesium sulphate, filtered and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a  
25 solution of two equivalents oxalic acid in absolute ethanol. The precipitate was dissolved in methanol, filtered, and concentrated under reduced pressure. the solid was recrystallised from absolute ethanol

**METHOD F:**

30 In a three-neck flask kept under nitrogen was placed a solution of the suitable phenol (1.6 mmol), 3-(diethylamino)propanol (1.5 mmol), and triphenyl phosphine (1.9 mmol) in 10 ml freshly distilled tetrahydrofuran. The mixture was stirred and cooled to 0°C with an ice and salt bath. A solution of diisopropyl

azodicarboxylate (2 mmol) in 10 ml tetrahydrofuran was added very slowly (typically over 40 minutes) and the mixture was allowed to warm to room temperature after which it was stirred overnight at room temperature (16 hours). The solvent was then removed under reduced pressure, the residue dissolved in ethyl acetate (20 ml) and the product extracted with 2N HCl (2x10 ml). The aqueous solution was neutralised with sodium hydroxide and the product extracted with dichloromethane. After drying over magnesium sulphate and filtration, the solvent was removed under reduced pressure. The residue was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol (compds. 43 and 46) or from a 1:1 mixture of methanol and absolute ethanol (compd. 42).

15

#### METHOD G:

A solution of the free base of compound 39 (0.6 mmol) or compound 46 (0.8 mmol) in 20 ml dry diethyl ether was added dropwise to a stirred suspension of lithium aluminium hydride (0.6 or 0.8 mmol) in 20 ml dry diethyl ether kept under nitrogen. The mixture was stirred at room temperature under nitrogen for two hours. Ice-cold water was carefully added and the organic layer decanted. The aqueous phase was extracted with diethyl ether. The combined organic solutions were dried over magnesium sulphate, filtered and concentrated under reduced pressure to leave a yellow oil. The oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. The precipitate was washed with diethyl ether and recrystallised from absolute ethanol (compd 50) or from isopropyl alcohol, giving a very hygroscopic solid (compd. 63).

#### METHOD H:

A solution of the suitable  $\alpha$ -bromo- $\omega$ -(4-cyanophenoxy) alkane (0.5 to 0.7 mmol) and an excess of the suitable secondary amine (8 to 12 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 24 hours

(compds. 54, 55, 57 and 60), 20 hours (compd. 52), 16 hours (compds. 56, 58, 59 and 61) or 8 hours (compd. 51) or was stirred at room temperature for 48 hours (compd. 53) or 24 hours (compd. 60). After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. Compound 62 was purified by column chromatography on silica gel eluting with ethyl acetate, and concentrated under reduced pressure. For all the compounds of method H, the remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol (two recrystallisations were required for compds. 58 and 59) or from a 1:1 mixture of methanol and absolute ethanol (compd. 55).

#### METHOD J:

A solution of compound 46 (1 mmol) in 10 ml methanol was stirred at room temperature and a solution of hydroxylamine hydrochloride (2 equivalents) in 2 ml water was added. The mixture was stirred at 50-70°C in a water bath for 20 minutes. Methanol was removed under reduced pressure. The residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. Compound 64 was purified by column chromatography on silica gel eluting with ethyl acetate, and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. Diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

**METHOD K:**

A solution of 4'-(3-bromopropoxy)acetophenone (0.8 to 1.9 mmol) and an excess of the suitable piperidine (3 to 10 equivalents) in 10 ml absolute ethanol was heated to reflux temperature with stirring for 16 hours. After cooling, the solvent was removed under reduced pressure and the residue diluted with aqueous sodium hydroxide. The product was extracted with diethyl ether, the organic extracts washed with water, dried over magnesium sulphate, filtered and concentrated under reduced pressure. The cis and trans isomers 67 and 70 were separated by column chromatography on silica gel eluting with a solvent mixture of diethyl ether, petroleum spirits (bp 60-80°C) and triethylamine in the ratio 66:33:1, and the eluent was removed under reduced pressure to leave an oil. Compound 75 was purified by column chromatography on silica gel eluting with chloroform and methanol (1:1), and concentrated under reduced pressure. The remaining oil was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents of oxalic acid in absolute ethanol. If no precipitate appeared, diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

**METHOD L:**

In a three-neck flask kept under nitrogen was placed a solution of the suitable 4'-hydroxyphenyl ketone (0.9 to 3 mmol), 3-(1-piperidiny)propanol (0.9 to 3 mmol), and triphenyl phosphine (1 to 3.5 mmol) in 10 ml freshly distilled tetrahydrofuran. The mixture was stirred and cooled to 0°C with an ice and salt bath. A solution of diethyl azodicarboxylate (1 to 3.6 mmol) in 10 ml tetrahydrofuran was added very slowly (typically over 40 minutes) and the mixture was allowed to warm to room temperature after which it was stirred overnight at room temperature (16 hours). The solvent was then removed under reduced pressure, the residue dissolved in ethyl acetate (20 ml) and the product extracted with 2N HCl (2x10 ml). The aqueous solution was neutralised with sodium hydroxide and the product extracted with dichloromethane. After drying over magnesium sulphate and filtration, the solvent was removed under reduced pressure. The crude product was purified by column chromatography

on silica gel eluting with diethyl ether containing 1% triethylamine, and concentrated under reduced pressure. The residue was converted to oxalate salt by dissolving in a small amount of absolute ethanol and adding a solution of two equivalents oxalic acid in absolute ethanol. If no precipitate appeared, 5 diethyl ether was added to form a precipitate. The solid was washed with diethyl ether and recrystallised from absolute ethanol.

### **Pharmacological study**

Interaction of compounds with the H<sub>3</sub> receptor are evidenced *in* 10 *vitro* by the measurement of the release of neosynthesized tritiated histamine from rat cerebral cortex synaptosomes preincubated with tritiated histidine (Garbarg et al., J. Pharmacol. Exp. Ther., 1992, 263 : 304-310). The H<sub>3</sub> potency of compounds is measured by the progressive reversal of the tritiated histamine release inhibition by the selective H<sub>3</sub> agonist (R) $\alpha$ -methylhistamine (Arrang et 15 al., Nature, 1987, 327: 117-123).

The effects of antagonists were estimated *in vivo* by the measurement of the tele-methylhistamine level variations in the brain of mice (Garbarg et al., J. Neurochem., 1989, 53: 1724-1730). At various time after p.o. administration of the compound, the effect of the H<sub>3</sub> antagonist is evidenced by 20 the increase in the telemethylhistamine level induced. This increase is compared to the maximal effect induced by the reference H<sub>3</sub>-antagonist thioperamide given at the dose of 10 mg/kg, p.o. This allows the calculation of the ED<sub>50</sub> value for each compound which correspond to the dose responsible for an half maximal effect.

25 The results are reported in the following table II:


Ex No.	X	n	R <sup>1</sup> R <sup>2</sup>	R <sup>3</sup> (n <sub>3</sub> = 1)	Ki (nM)	ED <sub>50</sub> (mg/kg/p.o.)
18	O	5	-(CH <sub>2</sub> ) <sub>4</sub> -	p-NO <sub>2</sub>	39 ± 11	1.1
43	O	3	Et, Et	p-CN	95 ± 28	0.50
46	O	3	Et, Et	p-CH <sub>3</sub> CO		0.44
50	O	5	-(CH <sub>2</sub> ) <sub>4</sub> -	p-CH <sub>3</sub> CH(OH)		1.0
56	O	4	Et, Et	p-CN		1.1
59	O	3	-(CH <sub>2</sub> ) <sub>5</sub> -	p-CN		0.20
60	O	3	-(CH <sub>2</sub> ) <sub>6</sub> -	p-CN		0.64
63	O	3	Et, Et	p-CH <sub>3</sub> CH(OH)		0.34
64	O	3	Et, Et	p-CH <sub>3</sub> C=N(OH)		0.8
66	O	3	-(3-Me)-(CH <sub>2</sub> ) <sub>5</sub> -	p-CH <sub>3</sub> CO		0.3
68	O	3	-(4-Me)-(CH <sub>2</sub> ) <sub>5</sub> -	p-CH <sub>3</sub> CO		0.3
69	O	3	-(CH <sub>2</sub> ) <sub>5</sub> -	p-C <sub>2</sub> H <sub>5</sub> CO		0.4

**THIS PAGE BLANK (USPTO)**




CLAIMS

1. Compound of general formula (I)




in which:

- $C_nH_{2n}$  is a linear or branched hydrocarbon chain with n ranging from 2 to 8;
  - X is an oxygen or sulfur atom;
  - $R^1$  and  $R^2$  may be identical or different and represent each independently
    - 10 • a lower alkyl or cycloalkyl,
- or taken together with the nitrogen atom to which they are attached,
- a saturated nitrogen-containing ring



with m ranging from 4 to 7, or

- an unsaturated nitrogen-containing ring



with p, q and r being 1 to 3 independently,

- 25 such nitrogen-containing ring i) or ii) being unsubstituted or substituted by one or more lower alkyl or cycloalkyl, or carboalkoxy groups, or
- a morpholino group, or
  - a N-substituted piperazino group:



with R being a lower alkyl, an alkanoyl or an optionally substituted phenyl group;

- $n_3$  is an integer from 0 to 5;

–  $R^3$  represents each independently

- a halogen atom,
- a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkyloximino, aryloximino,  $\alpha$ -hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group,
- or taken together with the carbon atoms of the phenyl ring to which it is fused, a 5- or 6-membered saturated or unsaturated ring or a benzene ring,

as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers, except compounds in which

- $-NR^1R^2$  is a pyrrolidinyl group,  $C_nH_{2n}$  is a linear chain  $-(CH_2)_n-$  and  $n_3$  is zero, X being an oxygen atom with n ranging from 3 to 5, or X being a sulfur atom with n being 4 or 5;
- $-NR^1R^2$  is a piperidinyl group,  $C_nH_{2n}$  is a linear chain  $-(CH_2)_n-$  and X is an oxygen atom,  $n_3$  being zero with n being 2, 5 or 8 or  $n_3$  being 1 with  $R^3$  being 4-CN and n being 5;
- $-NR^1R^2$  is a diethylamine group, X is an oxygen atom,  $C_nH_{2n}$  is a linear chain  $-(CH_2)_n-$  and  $n_3$  is 1,  $R^3$  being 4- $NO_2$  or 4- $COCH_3$  with n being 3 or  $R^3$  being 4-CN with n being 2 to 4;
- $-NR^1R^2$  is a dimethylamine group, X is an oxygen atom,  $C_nH_{2n}$  is a linear chain  $-(CH_2)_n-$  and  $n^3$  is 1,  $R^3$  being 4-CN with n being 3.

2. Compound according to claim 1, in which  $R^1$  and  $R^2$  are independently a lower alkyl group.

3. Compound according to claim 2, in which  $R^1$  and  $R^2$  are each an ethyl group.


4. Compound according to claim 1, in which  $-NR^1R^2$  is a saturated nitrogen-containing ring:



m being as defined in claim 1.

5        5. Compound according to claim 4, characterized in that m is 4, 5 or 6.

6. Compound according to claim 1, characterized in that  $-NR^1R^2$  is an unsaturated nitrogen-containing ring:



p, q and r being as defined in claim 1, preferably p, q and r are 1 or 2, more preferably p is 2 and q and r are 1.

15        7. Compound according to anyone of claims 4 to 6, characterized in that the nitrogen-containing ring i) or ii) is unsubstituted.

8. Compound according to anyone of claim 4 to 6, characterized in that the nitrogen-containing ring i) or ii) is substituted, preferably mono-substituted with an alkyl group.

20        9. Compound according to claim 8, characterized in that the nitrogen-containing ring is mono-substituted with a methyl group.

10. Compound according to claim 1, characterized in that  $-NR^1R^2$  is a morpholino group.

11. Compound according to claim 1, characterized in that  $-NR^1R^2$  is a N-substituted piperazino group, preferably N-acetylpiperazino.

25        12. Compound according to anyone of claims 1 to 11, characterized in that  $n_3$  is zero.

13. Compound according to anyone of claims 1 to 11, characterized in that  $n_3$  is 1 with  $R^3$  being as defined in claim 1 and preferably in para-position.

30        14. Compound according to anyone of claims 1 to 11 and 13, characterized in that  $R^3$  is a lower alkyl, preferably a  $C_1$ - $C_4$  alkyl.

15. Compound according to anyone of claims 1 to 11 and 13, characterized in that  $R^3$  is a halogen atom, a cyano, nitro, alkanoyl, alkyloximine

or hydroxyalkyl, preferably CN, NO<sub>2</sub>, COCH<sub>3</sub>, COC<sub>2</sub>H<sub>5</sub>, H<sub>3</sub>C-C=N-OH or H<sub>3</sub>C-CHOH.

16. Compound according to anyone of claims 1 to 11, characterized in that R<sup>3</sup> taken together with the carbon atoms of the phenyl group to which it is fused, form a 5- or 6- membered saturated or unsaturated ring, in particular a 5,6,7,8-tetrahydronaphthyl group.

17. Compound according to anyone of claims 1 to 11, characterized in that R<sup>3</sup> taken together with the phenyl group to which it is fused, form a naphthyl group.

18. Compound according to anyone of claims 1 to 17, characterized in that -C<sub>n</sub>H<sub>2n</sub>- is a linear hydrocarbon chain -(CH<sub>2</sub>)<sub>n</sub>-, n being as defined in claim 1.

19. Compound according to anyone of claims 1 to 18, characterized in that X is an oxygen atom.

20. Compound according to anyone of claims 1 to 18, characterized in that X is a sulfur atom.

21. Compound according to anyone of claims 1 to 20, characterized in that n is varying from 3 to 5 and is preferably 3.

22. Compound according to anyone of claims 1 to 21, characterized in that it is one of the following compounds:

N-methyl-N-(5-phenoxy-pentyl)-ethylamine

1-(5-phenoxy-pentyl)-morpholine

N-(5-phenoxy-pentyl)-hexamethyleneimine

N-ethyl-N-(5-phenoxy-pentyl)-propylamine

1-(5-phenoxy-pentyl)-2-methyl-piperidine

1-(5-phenoxy-pentyl)-4-propyl-piperidine

1-(5-phenoxy-pentyl)-4-methyl-piperidine

1-(5-phenoxy-pentyl)-3-methyl-piperidine

1-acetyl-4-(5-phenoxy-pentyl)-piperazine

1-(5-phenoxy-pentyl)-3,5-trans-dimethyl-piperidine

1-(5-phenoxy-pentyl)-3,5-cis-dimethyl-piperidine

1-(5-phenoxy-pentyl)-2,6-cis-dimethyl-piperidine

4-carboethoxy-1-(5-phenoxy-pentyl)-piperidine


	3-carboethoxy-1-(5-phenoxy-pentyl)-piperidine
	1-(5-phenoxy-pentyl)-1,2,3,6-tetrahydropyridine
	1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-chlorophenoxy)-pentyl]-pyrrolidine
5	1-[5-(4-methoxyphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-methylphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-cyanophenoxy)-pentyl]-pyrrolidine
	1-[5-(2-naphthyloxy)-pentyl]-pyrrolidine
	1-[5-(1-naphthyloxy)-pentyl]-pyrrolidine
10	1-[5-(3-chlorophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-phenylphenoxy)-pentyl]-pyrrolidine
	1-[5-[2-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine
	1-[5-(3-phenylphenoxy)-pentyl]-pyrrolidine
	1-(5-phenoxy-pentyl)-2,5-dihydropyrrole
15	1-[5-[1-(5,6,7,8-tetrahydronaphthyl)-oxy]-pentyl]-pyrrolidine
	1-(6-phenoxyhexyl)-pyrrolidine
	1-[5-(3-nitrophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-fluorophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-nitrophenoxy)-pentyl]-3-methyl-piperidine
20	1-[5-(4-acetylphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-aminophenoxy)-pentyl]-pyrrolidine
	1-[5-(3-cyanophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-benzoylphenoxy)-pentyl]-pyrrolidine
	1-[5-[4-(phenylacetyl)-phenoxy]-pentyl]-pyrrolidine
25	1-[5-(4-acetamidophenoxy)-pentyl]-pyrrolidine
	1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine
	1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine
	1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]-pyrrolidine
	1-[5-(4-cyanophenoxy)-pentyl]-diethylamine
30	N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine
	N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine
	1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine
	1-[3-(4-cyanophenoxy)-propyl]-piperidine

- N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine  
 N-[6-(4-cyanophenoxy)-hexyl]-diethylamine  
 N-[3-(4-cyanophenoxy)-propyl]-dipropylamine  
 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine  
 5 4-(3-diethylaminopropoxy)-acetophenone-oxime  
 1-[3-(4-acetylphenoxy)-propyl]-piperidine  
 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine  
 1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-piperidine  
 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine  
 10 1-[3-(4-propionylphenoxy)-propyl]-piperidine  
 1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine  
 1-[3-(4-formylphenoxy)-propyl]-piperidine  
 1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine  
 N-[3-(4-propionylphenoxy)-propyl]-diethylamine  
 15 1-[3-(4-butyrylphenoxy)-propyl]-piperidine  
 1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine  
 23. Compound according to anyone of claims 1 to 22,  
 characterized in that it is one of the following compounds :  
 1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine  
 20 1-[5-[4-(1-hydroxyethyl)-phenoxy]-pentyl]-pyrrolidine  
 1-[3-(4-cyanophenoxy)-propyl]-piperidine  
 N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine  
 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine  
 4-(3-diethylaminopropoxy)-acetophenone-oxime  
 25 1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine  
 1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine  
 1-[3-(4-propionylphenoxy)-propyl]-piperidine  
 24. Pharmaceutical composition characterized in that it  
 comprises as active ingredient, a therapeutically effective amount of a  
 30 compound according to anyone of claim 1 to 23 in combination with a  
 pharmaceutically acceptable vehicle or excipient.

25. Medicament acting as an antagonist of the histamine H<sub>3</sub>-receptors, characterized in that it comprises as active ingredient, an effective amount of a compound according to anyone of claims 1 to 23.

26. Use of a compound of general formula (I)


5



in which:

- C<sub>n</sub>H<sub>2n</sub> is a linear or branched hydrocarbon chain with n ranging from 2 to 8;
  - 10 - X is an oxygen or sulfur atom;
  - R<sup>1</sup> and R<sup>2</sup> may be identical or different and represent each independently
    - a lower alkyl or cycloalkyl,
- or taken together with the nitrogen atom to which they are attached,
- a saturated nitrogen-containing ring


15



with m ranging from 4 to 7, or

- an unsaturated nitrogen-containing ring


20



- 25 with p, q and r being 1 to 3 independently, such nitrogen-containing ring i) or ii) being unsubstituted or substituted by one or more lower alkyl or cycloalkyl, or carboalkoxy groups, or

- a morpholino group, or
- a N-substituted piperazino group:

30



with R being a lower alkyl, an alkanoyl or an optionally substituted phenyl group;

- n<sub>3</sub> is an integer from 0 to 5;

– R<sup>3</sup> represents each independently

- a halogen atom,
- a lower alkyl or cycloalkyl, a trifluoromethyl, aryl, alkoxy, aryloxy, nitro, formyl, alkanoyl, aroyl, arylalkanoyl, amino, carboxamido, cyano, alkyloximino, aryloximino,  $\alpha$ -hydroxyalkyl, alkenyl, alkynyl, sulphamido, sulfamoyl, carboxamide, carboalkoxy, arylalkyl or oxime group,
- or taken together with the carbon atoms of the phenyl ring to which it is fused, a 5- or 6-membered saturated or unsaturated ring or a benzene ring.

as well as their pharmaceutically acceptable salts, their hydrates, their hydrated salts, the polymorphic crystalline structures of these compounds and their optical isomers, racemates, diastereoisomers and enantiomers,

for the preparation of a medicament acting as an antagonist of the histamine H<sub>3</sub>-receptors.

27. Use according to claim 26, characterized in that compound (I) is as defined in any one of claims 2 to 21.

28. Use according to claim 26 characterized in that compound (I) is one of the following compounds:

- 1-(5-phenoxypropyl)-piperidine
- 1-(5-phenoxypropyl)-pyrrolidine
- N-methyl-N-(5-phenoxypropyl)-ethylamine
- 1-(5-phenoxypropyl)-morpholine
- N-(5-phenoxypropyl)-hexamethyleneimine
- N-ethyl-N-(5-phenoxypropyl)-propylamine
- 1-(5-phenoxypropyl)-2-methyl-piperidine
- 1-(5-phenoxypropyl)-4-propyl-piperidine
- 1-(5-phenoxypropyl)-4-methyl-piperidine
- 1-(5-phenoxypropyl)-3-methyl-piperidine
- 1-acetyl-4-(5-phenoxypropyl)-piperazine
- 1-(5-phenoxypropyl)-3,5-trans-dimethyl-piperidine
- 1-(5-phenoxypropyl)-3,5-cis-dimethyl-piperidine
- 1-(5-phenoxypropyl)-2,6-cis-dimethyl-piperidine



- 4-carboethoxy-1-(5-phenoxypropyl)-piperidine  
3-carboethoxy-1-(5-phenoxypropyl)-piperidine  
1-(5-phenoxypropyl)-1,2,3,6-tetrahydropyridine  
1-[5-(4-nitrophenoxy)-propyl]-pyrrolidine  
5 1-[5-(4-chlorophenoxy)-propyl]-pyrrolidine  
1-[5-(4-methoxyphenoxy)-propyl]-pyrrolidine  
1-[5-(4-methylphenoxy)-propyl]-pyrrolidine  
1-[5-(4-cyanophenoxy)-propyl]-pyrrolidine  
1-[5-(2-naphthoxy)-propyl]-pyrrolidine  
10 1-[5-(1-naphthoxy)-propyl]-pyrrolidine  
1-[5-(3-chlorophenoxy)-propyl]-pyrrolidine  
1-[5-(4-phenylphenoxy)-propyl]-pyrrolidine  
1-[5-{2-(5,6,7,8-tetrahydronaphthyl)-oxy}-propyl]-pyrrolidine  
1-[5-(3-phenylphenoxy)-propyl]-pyrrolidine  
15 1-(5-phenoxypropyl)-2,5-dihydropyrrole  
1-[5-{1-(5,6,7,8-tetrahydronaphthyl)-oxy}-propyl]-pyrrolidine  
1-(4-phenoxybutyl)-pyrrolidine  
1-(6-phenoxyhexyl)-pyrrolidine  
1-(5-phenylthiopropyl)-pyrrolidine  
20 1-(4-phenylthiobutyl)-pyrrolidine  
1-(3-phenoxypropyl)-pyrrolidine  
1-[5-(3-nitrophenoxy)-propyl]-pyrrolidine  
1-[5-(4-fluorophenoxy)-propyl]-pyrrolidine  
1-[5-(4-nitrophenoxy)-propyl]-3-methyl-piperidine  
25 1-[5-(4-acetylphenoxy)-propyl]-pyrrolidine  
1-[5-(4-aminophenoxy)-propyl]-pyrrolidine  
1-[5-(3-cyanophenoxy)-propyl]-pyrrolidine  
N-[3-(4-nitrophenoxy)-propyl]-diethylamine  
N-[3-(4-cyanophenoxy)-propyl]-diethylamine  
30 1-[5-(4-benzoylphenoxy)-propyl]-pyrrolidine  
1-[5-{4-(phenylacetyl)-phenoxy}-propyl]-pyrrolidine  
N-[3-(4-acetylphenoxy)-propyl]-diethylamine  
1-[5-(4-acetamidophenoxy)-propyl]-pyrrolidine

- 1-[5-(4-phenoxyphenoxy)-pentyl]-pyrrolidine  
1-[5-(4-N-benzamidophenoxy)-pentyl]-pyrrolidine  
1-[5-(4-(1-hydroxyethyl)-phenoxy)-pentyl]-pyrrolidine  
1-[5-(4-cyanophenoxy)-pentyl]-diethylamine  
5 1-[5-(4-cyanophenoxy)-pentyl]-piperidine  
N-[5-(4-cyanophenoxy)-pentyl]-dimethylamine  
N-[2-(4-cyanophenoxy)-ethyl]-diethylamine  
N-[3-(4-cyanophenoxy)-propyl]-dimethylamine  
N-[4-(4-cyanophenoxy)-butyl]-diethylamine  
10 N-[5-(4-cyanophenoxy)-pentyl]-dipropylamine  
1-[3-(4-cyanophenoxy)-propyl]-pyrrolidine  
1-[3-(4-cyanophenoxy)-propyl]-piperidine  
N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine  
N-[6-(4-cyanophenoxy)-hexyl]-diethylamine  
15 N-[3-(4-cyanophenoxy)-propyl]-dipropylamine  
N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine  
4-(3-diethylaminopropoxy)-acetophenone-oxime  
1-[3-(4-acetylphenoxy)-propyl]-piperidine  
1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine  
20 1-[3-(4-acetylphenoxy)-propyl]-3,5-trans-dimethyl-piperidine  
1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine  
1-[3-(4-propionylphenoxy)-propyl]-piperidine  
1-[3-(4-acetylphenoxy)-propyl]-3,5-cis-dimethyl-piperidine  
1-[3-(4-formylphenoxy)-propyl]-piperidine  
25 1-[3-(4-isobutyrylphenoxy)-propyl]-piperidine  
N-[3-(4-propionylphenoxy)-propyl]-diethylamine  
1-[3-(4-butyrylphenoxy)-propyl]-piperidine  
1-[3-(4-acetylphenoxy)-propyl]-1,2,3,6-tetrahydropyridine  
29. Use according to claim 26, characterized in that compound  
30 (I) is one of the following compounds:  
1-[5-(4-nitrophenoxy)-pentyl]-pyrrolidine  
N-[3-(4-cyanophenoxy)-propyl]-diethylamine  
N-[3-(4-acetylphenoxy)-propyl]-diethylamine

1-{5-[4-(1-hydroxyethyl)-phenoxy]-pentyl}-pyrrolidine  
N-[4-(4-cyanophenoxy)-butyl]-diethylamine  
1-[3-(4-cyanophenoxy)-propyl]-piperidine  
N-[3-(4-cyanophenoxy)-propyl]-hexamethyleneimine  
5 N-3-[4-(1-hydroxyethyl)-phenoxy]-propyl-diethylamine  
4-(3-diethylaminopropoxy)-acetophenone-oxime  
1-[3-(4-acetylphenoxy)-propyl]-3-methyl-piperidine  
1-[3-(4-acetylphenoxy)-propyl]-4-methyl-piperidine  
1-[3-(4-propionylphenoxy)-propyl]-piperidine

10 30. Medicament according to anyone of claims 25 to 29, for the treatment of central nervous system disorders, in particular Alzheimer disease, mood and attention alterations, cognitive deficits in psychiatric pathologies, obesity, vertigo and motion sickness.

15 31. Medicament according to anyone of claims 25 to 29, having psychotropic effects, promoting wakefulness, attention, memory and improving mood, intended to be used in particular in the treatment of Alzheimer disease and other cognitive disorders in aged persons, depressive or asthenic states.

20 32. Medicament according to anyone of claims 25 to 29, having nootropic effects, intended to be used in particular in treatment to stimulate attention and memorization capacity.

33. Medicament according to anyone of claims 25 to 29, for the treatment of obesity, vertigo and motion sickness.

34. Medicament according to anyone of claims 25 to 29, for the treatment of CNS disorders, in particular of aged persons.


**THIS PAGE BLANK (USPTO)**

Société Civile BIOPROJET

NON-IMIDAZOLE ARYLOXY (OR ARYLTHIO) ALKYLAMINES AS  
HISTAMINE H<sub>3</sub>-RECEPTOR ANTAGONISTS AND THEIR THERAPEUTIC  
APPLICATIONS.

ABSTRACT

Compounds of formula (I):



and their use for preparing medicaments acting as antagonists at the H<sub>3</sub>-receptors of histamine.

**THIS PAGE BLANK (USPTO)**